

# Numerical Study of the $S = 1$ Antiferromagnetic Spin Chain with Bond Alternation

Yusuke Kato and Akihiro Tanaka

*Department of Applied Physics, University of Tokyo, Bunkyo-ku, Tokyo 113, Japan*

(RECEIVED )

We study the  $S = 1$  quantum spin chain with bond alternation  $\mathcal{H} = \sum_i (1 - (-1)^i \delta) \mathbf{S}_i \cdot \mathbf{S}_{i+1}$  by the density matrix renormalization group method recently proposed by Steven R. White (Phys. Rev. Lett. **69** (1993) 3844). We find a massless point at  $\delta_c = 0.25 \pm 0.01$ . We also find the edge states in the region  $\delta < \delta_c$  under the open boundary condition, which disappear in the region  $\delta > \delta_c$ . At the massless point, the spin wave velocity  $v_s$  is  $3.66 \pm 0.10$  and the central charge  $c$  is  $1.0 \pm 0.15$ . Our results indicate that a continuous phase transition occurs at the massless point  $\delta = \delta_c$  accompanying breaking of the hidden  $Z_2 \times Z_2$  symmetry.

KEYWORDS: quantum antiferromagnetic spin chain, Haldane conjecture, bond alternation, density matrix renormalization group

The antiferromagnetic  $S = 1$  Heisenberg chain has been the subject of a large amount of interest in the last decade since Haldane argued the difference of the low energy physics between integer and half-integer spin [1,2]. The key concept of the Haldane conjecture is described in terms of the  $O(3)$  nonlinear sigma model with the topological term and he concluded that the topological term is ineffective and the spectrum is massive for integer spin, while the topological term suppresses the mass-generation and excitation spectrum is massless for half-integer spin. For the  $S = 1/2$  Heisenberg model, it is established that the excitation spectrum is gapless by the Bethe-Ansatz solution. The existence of excitation gap in the case of  $S = 1$  was verified from the theoretical [3,4], numerical [5–9], and experimental results [10–13].

Some years later, Haldane and Affleck [14–18] considered the antiferromagnetic (AF) spin chain with bond alternation  $\delta$ ,

$$\mathcal{H} = \sum_{i=1}^{N_s-1} \left(1 - (-1)^i \delta\right) \mathbf{S}_i \cdot \mathbf{S}_{i+1}. \quad (1)$$

When this Hamiltonian is mapped to the continuum model, *i.e.*, the  $O(3)$  nonlinear sigma model, the topological angle  $\theta$ , which is the coefficient of the Berry phase term in the action, is given by  $\theta = 2\pi S(1 - \delta)$ . According to the argument by Haldane and Affleck, only in the case where  $\theta = \pi \times (\text{odd integer})$ , is the system in the critical state with massless excitations. Hence as the bond alternation  $\delta$  changes from  $-1$  to  $1$ , the system experiences critical (massless) point  $2S$  times. Applied to the  $S = 1/2$  case, this general discussion is consistent with Cross-Fisher’s argument of the spin-Peierls transition [19]. However the situation is nontrivial in the case of  $S = 1$ . Without the bond alternation  $\delta$ , the system is gapful, *i.e.*, possesses the Haldane gap, and the above discussion predicts that the massless point appears at some  $\delta = \delta_c \neq 0$ . However this argument contains several approximations and assumptions, such as taking the large  $S$  limit, and it is important to study the model (1) by another approach and to give an independent check of the above scenario.

Varying- $\delta$  problem in the model (1) can be attacked from the viewpoint of the topological order proposed by den-Nijs and Rommelse [20] and Tasaki [21]. This “hidden” order is

measured by the string correlation function

$$g(i, j) = \langle S_i^z \left( \prod_{k=i+1}^{j-1} e^{i\pi S_k^z} \right) S_j^z \rangle. \quad (2)$$

It is widely believed that this correlation function is long-ranged at the Heisenberg point  $\delta = 0$  from numerical evidences [22,23]. On the other hand, an elementary calculation shows that the string correlation function is exactly zero at the completely dimerized points  $\delta = \pm 1$ . Hence we expect that phase transitions corresponding to the hidden symmetry breaking should occur twice; once in the interval between  $\delta = -1$  and  $\delta = 0$  and once more between  $\delta = 0$  and  $\delta = 1$ . However, these arguments cannot determine whether the phase transitions are discontinuous or continuous. Thus the existence of the massless points as predicted by Haldane and Affleck is still an open problem and direct numerical study is needed. In this paper we investigate the model (1) with  $S = 1$  numerically for the region  $0 \leq \delta \leq 1$ , using exact diagonalization and the density matrix renormalization group (DMRG) method. This is sufficient since reversing the sign of  $\delta$  merely corresponds to interchanging the roles of the two sublattices, and hence has no effects on the bulk properties. It does however have an intrinsic boundary effect for open chains on which we will remark later.

DMRG, which was recently proposed by White, is a new algorithm for real-space renormalization group [23–26]. The key point lays in the method of the truncation procedure of the Hilbert space. In conventional real space renormalization group calculation, one keeps the lowest-lying eigenstates of the block Hamiltonian in forming a new effective Hamiltonian. In the DMRG method, on the other hand, one keeps the most important eigenstates of the block density matrix, obtained from diagonalizing the Hamiltonian of a larger section of the lattice which includes the block. The above change brings high accuracy and the new algorithm is already applied to the  $S=1$  antiferromagnetic Heisenberg spin chain successfully [23–26]. For details of the DMRG algorithm, readers are referred to ref. [26].

Using the DMRG method, we calculated the excitation gap of systems of large length up to about two hundred sites and extrapolated them to the thermodynamic limit. The

numbers of state kept per block in our calculations are 27, 36, 48, and 81. As a benchmark test we calculated the ground state energy density of  $S = 1/2$  Heisenberg model. Comparing with the exact Bethe-Ansatz solution, we find that the relative error is about  $5 \times 10^{-4}$ . We also compared those of DMRG with that of exact diagonalization in the case of  $S = 1$  and site number  $N_s = 12$ . The two results were found to agree well within a relative error of  $7 \times 10^{-6}$ .

Figure 1 shows the excitation spectrum of  $S = 1$  spin chain versus dimer strength. The data of diagonalization (points plotted with triangles, squares, empty and filled circles and empty and filled diamonds) shows the gap between the singlet ground state and the first excited (triplet) state under periodic boundary condition. The results of diagonalization show that the gap has a minimum around  $\delta \sim 0.25$ , which has been already reported in ref. [27]. The minimum value of the gap becomes smaller with increasing system size. In the large  $N_s$  limit, there are two possibilities *i.e.* the minimum value of the gap tends to zero or saturates at a finite value. Limited system sizes in the diagonalization, however, disallows us to conclude which is realized in the thermodynamic limit.

The data of DMRG calculation (represented by the double circles and crosses), which are extrapolated values ( $N_s \rightarrow +\infty$ ), show the gap between the ground state and the first excited state, and that between the ground state and the second excited state under the open boundary condition. The gap behaviors are different between the region  $\delta < 0.25$  and  $\delta > 0.25$ . In the region  $\delta < 0.25$ , the first excited state is triplet ( $S_{\text{tot}} = 1$ ) and the second one is quintuplet ( $S_{\text{tot}} = 2$ ) while both states are triplet in the region  $\delta > 0.25$ . The first excited state in the region  $\delta < 0.25$ , approaches the singlet ground state exponentially with increasing system size. These triplet excitations have been investigated at the Heisenberg point ( $\delta = 0$ ) and are known to be the edge excitations near the open boundary of the spin chain [28,29]. For the moment, we focus upon the gap between ground state and the second excited state in the region  $\delta < 0.25$ , because the above “edge excitations” seem to have little effects upon the bulk properties of the spin chain. Later we will discuss the edge states, especially their relevance to the hidden symmetry breaking.

The size dependence of the gap between the ground state and the second excited state in the region  $\delta < 0.25$  is fitted well with  $1/N_s^2$ . The gap between the ground state and the first and second excited states in the region  $\delta > 0.25$  is also fitted well with  $1/N_s^2$ . At  $\delta = 0.25$ , size dependence of the gap behaves as  $1/N_s$ . In contrast to the exact diagonalization method, DMRG gives a clear evidence for the existence of a massless point at  $\delta_c = 0.25 \pm 0.01$ .

We now turn to the critical properties of the massless point. It is plausible to assume that at the massless point, the system has conformal invariance with respect to the low energy properties. Actually, we observe that the size dependence of the excitation gap is proportional to  $1/N_s$ , which is consistent with the finite size scaling of the conformal field theory (CFT). CFT gives the expression for the size dependence of ground state energy [30,31];

$$\frac{E_{g,s}(N_s)}{N_s} = \epsilon + \frac{f}{N_s} - \frac{\pi c v_s}{24 N_s^2} + \mathcal{O}\left(\frac{1}{N_s^3}\right), \quad (3)$$

under the open boundary condition. Here  $E_{g,s}(N_s)$ ,  $\epsilon$ ,  $f$ , and  $c$  are the ground state energy of finite size system, ground state energy density of the infinite system, surface energy and the central charge. We estimate  $v_s = 3.66 \pm 0.10$  at  $\delta = \delta_c$  from the excitation gap. Combining the value of  $v_s$  with the results for the ground state energy, we estimate  $c = 1.0 \pm 0.15$ .

Affleck and Haldane predicted  $\delta_c = 0.5$ , which deviates from our result  $\delta_c = 0.25 \pm 0.01$ . We can attribute this discrepancy to the mapping of the quantum spin chain onto the  $O(3)$  sigma model, which is justified in the large  $S$  limit. However their prediction is correct with regards to the *existence* of a massless point.

The existence of edge states or four-fold degeneracy of the ground state in the open chain was discussed theoretically [3,4,28,32] and was related to the long range topological order and hidden  $Z_2 \times Z_2$  symmetry breaking. So we conclude that at the massless point a continuous phase transition with the hidden  $Z_2 \times Z_2$  symmetry breaking occurs. Our result ( $c \sim 1$ ) is in accord with the general belief concerning the  $Z_2 \times Z_2$  symmetry breaking [33] that the massless phase boundary which corresponds to a full  $Z_2 \times Z_2$  symmetry breaking belongs to the same universality class as the Gaussian model.

We note in passing the consequence of our choice of the sign of  $\delta$ ; had we taken negative  $\delta$ , we would have found that the ground state degeneracy is nine-fold in the region  $-1 \leq \delta \leq \delta_c^-$  (due to the formation of two localized  $S = 1$  objects at the open ends), while it is still four-fold in the region  $\delta_c^- \leq \delta < 0$ , where  $\delta_c^- \equiv -\delta_c \sim -0.25$ .

In summary, we presented numerical results on the  $S = 1$  antiferromagnetic spin chain with bond alternation and concluded that the continuous phase transition occurs at  $\delta_c = 0.25 \pm 0.01$  with the hidden  $Z_2 \times Z_2$  symmetry breaking. The higher  $S$  cases are left for future investigations.

We are grateful to Professor N. Nagaosa for introducing us to this problem and for continuous encouragement as well as for critical reading of the manuscript. This work is supported by a Grant-in-Aid for Scientific Research on Priority Areas, “Computational Physics as a New Frontier in Condensed Matter Research” from the Ministry of Education, Science, and Culture of Japan. Numerical calculations were performed on HITAC S-3800 at the computer center of University of Tokyo. We also acknowledge Professor M. Kaburagi, Professor T. Tonegawa, and Dr. T. Nishino for their diagonalization program KOBEPACK/1.

## REFERENCES

- [1] F. D. M. Haldane: Phys. Lett. **93A** (1983) 464.
- [2] F. D. M. Haldane: Phys. Rev. Lett. **50** (1983) 1153.
- [3] I. Affleck, T. Kennedy, E. H. Lieb, and H. Tasaki: Phys. Rev. Lett. **59** (1987) 799.
- [4] I. Affleck, T. Kennedy, E. H. Lieb, and H. Tasaki: Commun. Math. Phys. **115** (1988) 477.
- [5] R. Botet and R. Jullien: Phys. Rev. B **27** (1983) 613.
- [6] R. Botet, R. Jullien, and M. Kolb: Phys. Rev. B **28** (1983) 3914.
- [7] M. Kolb, R. Botet, R. Jullien: J. Phys. **A16** (1983) L673.
- [8] M. P. Nightingale and H. W. Blöte: Phys. Rev. B **33** (1986) 659.
- [9] T. Sakai and M. Takahashi: Phys. Rev. B **42** (1990) 1090.
- [10] J. P. Renard, M. Verdaguer, L. P. Regnault, W. A. C. Erkelens, J. Rossat-Mignod, and W. G. Stirling: Europhys. Lett. **3** (1987) 945.
- [11] J. P. Renard, M. Verdaguer, L. P. Regnault, W. A. C. Erkelens, J. Rossat-Mignod, J. Ribas, W. G. Stirling, and C. Vettier: J. Appl. Phys. **63** (1988) 3538.
- [12] K. Katsumata, H. Hori, T. Takeuchi, M. Date, and A. Yamagishi, and J. P. Renard: Phys. Rev. Lett. **63** (1989) 86.
- [13] Y. Ajiro, T. Goto, H. Kikuchi, T. Sakakibara, and T. Inami: Phys. Rev. Lett. **63** (1989) 1424.
- [14] F. D. M. Haldane: J. Appl. Phys. **57** (1985) 3359.
- [15] I. Affleck: Phys. Rev. Lett. **54** (1985) 966.
- [16] I. Affleck: Nucl. Phys. **B257** (1985) 397.

- [17] I. Affleck: Nucl. Phys. **B265** (1986) 409.
- [18] I. Affleck and F. D. M. Haldane: Phys. Rev. B **36** (1987) 5291.
- [19] M. C. Cross and D. S. Fisher: Phys. Rev. B **19** (1979) 402.
- [20] M. den Nijs and K. Rommelse: Phys. Rev. B **40** (1989) 4709.
- [21] H. Tasaki: Phys. Rev. Lett. **66** (1991) 798.
- [22] S. M. Girvin and D. P. Arovas: Phys. Scr. **T27** (1989) 156.
- [23] S. R. White and D. A. Huse: Phys. Rev. B **48** (1993) 3844.
- [24] S. R. White: Phys. Rev. Lett. **69** (1992) 2863.
- [25] E. S. Sorensen and I. Affleck: Phys. Rev. Lett. **69** (1992) 2863.
- [26] S. R. White: Phys. Rev. B **48** (1993) 10345.
- [27] D. Guo, T. Kennedy, and S. Mazumdar: Phys. Rev. B **41** (1990) 9592.
- [28] T. Kennedy: J. Phys.:Condens. Matter **2** (1990) 5737.
- [29] M. Hagiwara, K. Katsumata, I. Affleck, and B. I. Halperin: Phys. Rev. Lett. **65** (1990) 3181.
- [30] H. W. J. Blöte, J. L. Cardy, and M. P. Nightingale: Phys. Rev. Lett. **56** (1986) 742.
- [31] I. Affleck: Phys. Rev. Lett. **56** (1986) 746.
- [32] T. Kennedy and H. Tasaki: Phys. Rev. B **45** (1992) 304.
- [33] This is established on rigorous grounds for the Ashkin-Teller model; L. Kadanoff and A. Brown: Ann. Phys. **121** (1979) 318.



## FIGURES

FIG. 1. Excitation gap between the ground state and excited states versus the dimer strength  $\delta$ . We show here two kinds of numerical results. Points plotted with triangles, squares, circles (empty and filled) and diamonds (empty and filled) are the gap between the ground state and the first excited state obtained by the exact diagonalization under the periodic boundary condition for system sizes  $N_s = 4, 6, 8, 10, 12$ , and  $14$ . As the system becomes larger, the minimum of the gap shows up around  $\delta \sim 0.25$ . Double circles/crosses represent the gap between the ground state and the first/second excited state obtained by the DMRG method (the infinite lattice method) under the open boundary condition. We find a massless point around  $\delta_c = 0.25 \pm 0.01$ , and also edge states (Kennedy triplet) in the region  $\delta < \delta_c$ .